

Welke depressieve patiënten hebben baat bij het toevoegen van kortdurende psychodynamische therapie aan antidepressiva?

dr. Ellen Driessen

ellen.driessen@ru.nl

NedKAD conferentie 2023

Expertisecentrum Depressie



Pro Persona
geestelijke gezondheidszorg

Welke depressieve patiënten hebben baat bij het toevoegen van kortdurende psychodynamische therapie aan antidepressiva?

Disclosure belangen spreker


Sponsoring/onderzoeksgeld

- American Psychoanalytic Association
- Stichting Psychoanalytische Fondsen
- NWO







A.




>




B.




||




C.




>



B.



=





Post-treatment: Cohen's $d = 0.26$

Follow-up: Cohen's $d = 0.50$



+



Driessen et al., 2020. *Clinical Psychology Review* <https://doi.org/10.1016/j.cpr.2020.101886>

Meta-analyse van individuele patiëntdata (IPD)



Original Article

Cite this article: Driessen E *et al* (2022). Which patients benefit from adding short-term psychodynamic psychotherapy to antidepressants in the treatment of depression? A systematic review and meta-analysis of individual participant data. *Psychological Medicine* 1–12. <https://doi.org/10.1017/S0033291722003270>

Received: 2 May 2022

Revised: 29 September 2022

Accepted: 30 September 2022

Key words:













Combined treatment; depression; individual participant data meta-analysis; moderator; short-term psychodynamic psychotherapy (STPP)

Author for correspondence:

Ellen Driessen,

E-mail: ellen.driessen@ru.nl

Which patients benefit from adding short-term psychodynamic psychotherapy to antidepressants in the treatment of depression? A systematic review and meta-analysis of individual participant data

Ellen Driessen^{1,2,3} , Marjolein Fokkema⁴ , Jack J. M. Dekker⁵ ,
Jaap Peen⁵ , Henricus L. Van⁶ , Giuseppe Maina^{7,8} , Gianluca Rosso^{7,8} ,
Sylvia Rigardetto⁸, Francesco Cuniberti⁷ , Veronica G. Vitriol⁹ ,
Antonio Andreoli¹⁰, Yvonne Burnand¹⁰, Jaime López Rodríguez¹¹,
Valerio Villamil Salcedo¹¹ , Jos W. R. Twisk¹², Frederik J. Wienicke¹ 
and Pim Cuijpers³ 

¹Department of Clinical Psychology, Behavioural Science Institute, Radboud University, Nijmegen, Netherlands; ²Depression Expertise Centre, Pro Persona Mental Health Care, Nijmegen, Netherlands; ³Department of Clinical, Neuro and Developmental Psychology, Amsterdam Public Health Research Institute, Vrije Universiteit Amsterdam, Amsterdam, Netherlands; ⁴Department of Methodology and Statistics, Leiden University, Leiden, Netherlands; ⁵Department of Research, Arkin Mental Health Care, Amsterdam, Netherlands; ⁶NPI, Arkin Mental Health Care, Amsterdam, Netherlands; ⁷Department of Neuroscience ‘Rita Levi Montalcini’, University of Turin, Turin, Italy; ⁸Psychiatric Unit, San Luigi Gonzaga University Hospital of Orbassano, Turin, Italy; ⁹Medical School, University of Talca, Talca, Chile; ¹⁰Geneva University Hospital Center, Geneva, Switzerland; ¹¹National Institute of Psychiatry, Mexico City, Mexico and ¹²Department of Epidemiology and Data Science, Amsterdam University Medical Centers,

Ernstigere depressieve klachten

Langere duur depressieve episode



+





Bedankt!

ellen.driessen@ru.nl